

A-T Clinical Center
Johns Hopkins Children's Center
Jenny Wright, RN - Clinic Coordinator
410.614.1922
jenny.wright@jhmi.edu

Howard M. Lederman, MD PhD
Clinic Director
410.955.5883
hlederm1@jhmi.edu

A-T Cancer Consultations
St. Jude Children's Research Hospital
John T. Sandlund, MD
901.495.3300
(to be contacted only by physicians per St. Jude policy)

Management of Acute Infection

A-T patients are predisposed to upper and lower respiratory tract infections because of immunodeficiency, aspiration, and impaired cough.

- Use of antibiotics should be considered for treatment of upper or lower respiratory tract infections that are severe, accompanied by fever, or persist for greater than 7 days.
- Precautions should be taken to reduce the risk of aspiration during respiratory illnesses.

Preoperative Evaluation and Surgery

A preoperative evaluation of lung function should be performed in all A-T patients regardless of age and whether or not they have chronic respiratory symptoms.

- Children and adults with A-T may have difficulty coming off the ventilator after surgery or other procedures requiring general anesthesia.
- Possible alternatives to general anesthesia and strategies that maximize airway clearance following anesthesia should be considered.
- Following placement of a gastrostomy tube, enteral feedings should be advanced with caution.



atcp.org | Tel: 800.5.HELP.A-T or 954.481.6611 | info@atcp.org

5300 W. Hillsboro Blvd., Suite 105, Coconut Creek, FL 33073 USA

QUICK INFORMATION FOR THE CARE OF PATIENTS WITH ATAXIA-TELANGIECTASIA

NAME

DATE OF BIRTH

Ataxia-telangiectasia (A-T) is an autosomal recessive disease characterized by:

- progressive neurologic degeneration with ataxia, eye movement abnormalities, dysarthria and impaired swallowing,
- immunodeficiency with lymphopenia and hypogammaglobulinemia of varying severity,
 - I have antibody deficiency and receive gammaglobulin therapy
- predisposition to a wide variety of cancers (lymphoma and leukemia most common, especially in children),
- telangiectasia (often present), especially over the sclerae. In rare instances, telangiectasia may cause complications in bladder, brain, other organs.

Diagnostic X-Rays and CT Scans in A-T

People with A-T have an increased sensitivity to ionizing radiation (x-rays and gamma rays).

- X-rays should be performed only when the result will affect medical management.
- If a patient with A-T has fever, cough, and breath sounds characteristic of pneumonia, a diagnosis of pneumonia can be made clinically and antibiotics can be prescribed without x-ray confirmation. If symptoms persist despite antibiotics, a follow-up chest x-ray may be useful.
- Routine screening dental x-rays should be avoided, but an x-ray to evaluate tooth pain is reasonable.
- In order to keep radiation to a minimum, patients should receive frontal view chest radiographs or radiation-sparing techniques for CT (e.g. fewer CT images per scan).
- There is no contraindication to MRI or ultrasound investigation.